

D. Schwartz · H. de Foresta · A. Mariotti
J. Balesdent · J.P. Massimba · C. Girardin

Present dynamics of the savanna-forest boundary in the Congolese Mayombe: a pedological, botanical and isotopic (^{13}C and ^{14}C) study

Received: 16 February 1995 / Accepted: 25 November 1995

Abstract Isolated savannas enclosed by forest are especially abundant in the eastern part of the Congolese Mayombe. They are about 3000 years old, and were more extensive some centuries ago. The boundary between forest and savanna is very abrupt, as a consequence of the numerous savanna fires lit by hunters. Floristic composition and vegetation structure data, organic carbon ratios, $\Delta^{14}\text{C}$ and $\delta^{13}\text{C}$ measurements presented here show that forest is spreading over savanna at the present time and suggest that the rate of forest encroachment is currently between 14 and 75 m per century, and more probably about 20–50 m per century. As most savannas are less than 1 km across, such rates mean, assuming there are no changes in environmental conditions, that enclosed savannas could completely disappear in the Mayombe in about 1000–2000 years.

Key words Congo · Forest-savanna dynamics · ^{13}C natural abundance · Forest encroachment · Pioneer trees

Introduction

All over the African equatorial forest, but especially in the Congolese Mayombe, forest cover encloses some isolated savannas. It is widely held that the enclosed savannas of the Mayombe are recent anthropic savannas. This hypothesis has been supported only by the fact that these savannas were densely occupied by villages at the beginning of the twentieth century (Vennetier 1968; Cusset 1989; Gibert and Sénéchal 1989; Petit 1990). As demonstrated by Foresta (1990), savanna formation due to man's activities may be, in the Mayombe, unfounded.

D. Schwartz (✉)¹ · H. de Foresta² · J.P. Massimba
ORSTOM, B.P. 1286, Pointe Noire, Congo

A. Mariotti · J. Balesdent · C. Girardin
Université P. et M. Curie – INRA, Biogéochimie Isotopique,
Case n° 120, 4 Place Jussieu, F-75252 Paris 5, France

Present addresses:

¹ CEREG, 3 rue de l'Argonne, F-67083 Strasbourg, France
² ICRAF, P.O. Box 161, Bogor 16001, Indonesia

Despite the availability of efficient tools such as the chain saw, the development of commercial agriculture, and the fact that vegetation in the most inhabited areas testifies to an important forest degradation resulting from repeated agricultural clearing and that villages are located in the less humid areas of the Mayombe, savanna formation can never be observed. Data collected suggest that these savannas have a palaeoclimatic origin (Foresta 1990; Schwartz et al. 1990b), that they could be 3000 years old and were more extensive during past centuries. Moreover, archaeological data suggest that they were previously found as far as the Les Saras area, which is currently purely forested (Schwartz et al. 1990a). This evidence suggests that forest is at present encroaching onto savanna. Botanical observations, both floristic and structural, agree with this hypothesis (Massimba 1987; Foresta 1990), but this approach is only qualitative. To provide more conclusive evidence, other approaches have to be used, such as determination of the mean residence time (MRT) of soil organic matter (SOM) by ^{14}C dating combined with an isotopic (^{13}C) characterization of vegetation change recorded in SOM. This paper presents the results from a combined botanical and SOM approach along two forest-enclosed savanna sequences, as a means to understand the present dynamics of the vegetation in the Congolese Mayombe.

Materials and methods

Study sites

The location of the observations is shown in Figs. 1 and 2. The Mayombe is a mountain extending from Gabon to Zaïre parallel to the coastline. In the Congo, the highest crests culminate at about 900 m. The rainfall increases from the west (1200 mm/year) to the highest crests (1980 mm/year) and then decreases down to Loubomo (1160 mm/year). Rain falls from October until May. Mists and drizzles are abundant during the dry season, and contribute to the retention of some moisture in soils. The number of hours of sunshine is small, especially during the dry season: less than 1000 h/year at Dimonika (Clairac et al. 1989). This dry, fresh and cloudy weather explains the presence of rain-forest in the less humid areas. The soils of the Mayombe are ferralic cambisols (FAO




Fig. 1 The Congolese Mayombe (1 coastal savannas, 2 forest-savanna mosaic of the western Congo, 3 forest areas, 4 savannas of the Niari-Nyanga synclinorium). The inset corresponds to Fig. 2




Fig. 2 Localization of the intramayombian savannas (after Schwartz et al. 1990b); (1 Forest areas, 2 intramayombian savannas, 3 savannas of the Niari valley)

1976). A stone-line is present in about 80–90% of the soils. Its thickness ranges between 5 cm and 2 m. It is located just above the weathering horizons. The horizons above the stone-line form a cover the thickness of which ranges between 0 and 5–6 m, and more often between 0.5 and 2 m. These variations are linked to the erosion-colluviation processes (Gras 1970). The vegetation is composed essentially of rain-forest of an “evergreen transitional to a semi-deciduous” type (White 1983). The forest cover of the Mayombe includes isolated enclosed savannas. They are more numerous in the eastern Mayombe, where villages are rare and human influence is weak (Vennetier 1968; Schwartz et al. 1990b). A few are found right in the middle of the forest (Fig. 2).

The outline of savannas is geometric and angular. The boundary with the forest is as a general rule abrupt (cf. infra), which permits the definition without ambiguity of the limit between the two

kinds of vegetation. In the savannas located north of Makaba, the main grass species is *Pobeguinea arrecta* (Gramineae). Occasional shrubs belong to familiar species: *Annona senegalensis*, *Bridelia ferruginea*, and *Nauclea latifolia*. It should be noticed that these savannas are burnt almost every year, even in the areas farthest from the villages. These bushfires are set by hunters to help their progression between forest zones.

Methods

Two forest-enclosed savanna transects are detailed. The first one is located to the WNW of Makaba, the second one to the west of Kwilila (Fig. 2). They are respectively 47 and 71 m long. The transects were carried out parallel to the contour lines in order to be as free as possible from erosion and colluviation processes. Makaba-transect was completed by a vegetation profile, carried out on $20 \times 50 = 1000 \text{ m}^2$. As the vegetation profile was 20 m wide, and isotopic transects were linear, some differences appear in the vegetation limits as shown in Figs. 3–5. Only the forest part of the vegetation profile was studied, because the savanna vegetation is mainly composed of *P. arrecta* (Gramineae). It is possible to distinguish, in a simplified manner, four zones in the transects: in succession (1) a graminean zone (savanna *stricto sensu*), (2) an herbaceous zone dominated by ferns, (3) the edge of the forest which forms a thicket composed of pioneer trees and above all shrubs, and (4) the forest. The savanna-forest interface corresponds to the limit between ferns and shrubs from a botanical viewpoint, whereas the isotopic interface corresponds to the C3–C4 vegetation change, i.e. the limit between Gramineae and ferns. We defined the zero point of distances along the transects as the abrupt Gramineae-fern contact, as a consequence of the C3–C4 vegetation changes. Analytical data were derived from the determination of organic carbon content of soils, by measurement of the $\delta^{13}\text{C}$ and ^{14}C datings of SOM.

Soil samples were collected from 10 cm \times 10 cm plots between the surface of the soil and a depth of 5 cm. Carbon content and ^{13}C natural abundance were determined on each sample, whereas the ^{14}C datings were carried out on several bulked samples collected at different points (savanna, edge, forest) of the transects. Soil organic carbon content was determined by the Walkley and Black (1934) method, which is suitable for ferrallitic soils.

The method of measurement of the ^{13}C natural abundance is given in detail by Girardin and Mariotti (1991). The ^{13}C content of a sample is expressed in delta units:


$$\delta^{13}\text{C} \text{ ‰} = [(^{13}\text{R sample} - ^{13}\text{R standard}) / ^{13}\text{R standard}] \times 1000$$

where ^{13}R stands for the isotopic ratio $^{13}\text{C}/^{12}\text{C}$. Data are expressed versus the PDB standard.

The standard deviation, as determined by 40 measurements on a well-homogenized soil sample, is $\pm 0.07 \text{ ‰} \delta^{13}\text{C}$. The ^{13}C content of plants varies according to their photosynthetic pathway (Bender 1971; Smith and Epstein 1971). Herbaceous plants of cold and temperate regions and all trees are C3 plants (Calvin cycle). They contain less ^{13}C than C4 plants (Hatch-Slack cycle), which are essentially tropical Gramineae and Cyperaceae. The $\delta^{13}\text{C}$ values of C3 plants are mostly between -23 and -34 ‰ (average of -26 ‰) whereas C4 plants range from -10 to -17 ‰ (average of -12 ‰).

Mineralization of SOM and pedogenetic processes which occur during humus evolution lead to only slight variation in the ^{13}C ratio observed with depth in the soil profile (Troughton et al. 1974; Goh et al. 1976, 1977; O'Brien and Stout 1978; Dzurec et al. 1985; Nadelhoffer and Fry 1988; Martin et al. 1990; Mariotti 1991; Balesdent et al. 1993; Mariotti and Peterschmitt 1994). Consequently, the organic matter $\delta^{13}\text{C}$ in a soil in equilibrium with vegetation of a precise photosynthetic pathway is very similar to that in the dominant vegetation. Schwartz (1991) and Schwartz et al. (1986) show that it is possible in the Congo to differentiate without doubt forest humus ($\delta^{13}\text{C} \approx -26.5$ to -28 ‰) from savanna humus, where Gramineae are widely dominant ($\delta^{13}\text{C} \approx -12$ to

Fig. 3 Schematic profile of the vegetation for transect no 1, Makaba. Parcel of 20×50 m=1000 m², except for 1 cm <DBH <10 cm trees (=5×50 m). Tree serial numbers correspond to Table 2



-16 ‰). At the very most, it is possible to notice a slight increase of several ‰ with depth in the soil (Balesdent et al. 1987; Volkoff and Cerri 1987; Mariotti and Peterschmitt 1994). This enrichment can be explained by different factors: isotopic fractionation during the degradation of organic matter by micro-organisms, corresponding to the emission of ¹³C depleted CO₂ with a correlative enrichment of the remaining organic matter (Kaplan and Rittenberg 1964; Blair et al. 1985); differential preservation of ¹³C enriched SOM components (Nissenbaum and Schallinger 1974; Benner et al. 1987); a decrease of around 1.5 ‰ in the atmospheric CO₂ δ¹³C during the last 150 years, as a consequence of the emission into the atmosphere of CO₂ resulting from the combustion of fossil fuels (Suess effect) and also from an increase in the SOM mineralization after cultivation (Marino and McElroy 1991). The δ¹³C from these two CO₂ sources is close to -25 ‰, which is very different from the atmospheric CO₂ (-7 to -8 ‰). Although SOM becomes ¹³C-enriched with depth in profiles, this enrichment (<4 ‰) is not large enough to mask the 14 ‰ difference between C3 and C4 plant litter. So, any vegetation change between C3 and C4 plants introduces a natural in situ labelling of organic matter

incorporated into the soil (Flexor and Volkoff 1977, Schwartz et al. 1986; Balesdent et al. 1987; Mariotti and Balesdent 1990; Martin et al. 1990).

The ¹⁴C datings were performed in the Radiocarbon Laboratory of the Ecole de Géologie de Nancy on humus samples by the classical liquid scintillation counting method. Radiocarbon activities were corrected for isotope fractionation and expressed as Δ¹⁴C units, according to Craig's equation:

$$\Delta^{14}\text{C} = \delta^{14}\text{C} - (2\delta^{13}\text{C} + 50)(1 + \delta^{14}\text{C}/1000).$$

Positive δ¹⁴C values reflect the incorporation of thermonuclear bomb radiocarbon in SOM, following the thermonuclear tests in the 1960s (O'Brien and Stout 1978). We used this natural tracer to calculate the mean residence time of SOM according to Balesdent and Guillet (1982). To compute the MRT-Δ¹⁴C relationship, we integrated the radiocarbon activity of the atmospheric CO₂ incorporated in the soil until the date of sampling (e.g. 1990) with the assumption that SOM is a well-mixed compartment with an exponential decay. The datings were carried out in three zones of each




Fig. 4 Organic carbon contents in the 0–5 cm horizon for Makaba and Kwilila transects




Fig. 5 $\delta^{13}\text{C}$ values of soil organic matter in the 0–5 cm horizon for Makaba and Kwilila transects, and mean residence time values in each vegetation zone (mean sample obtained by gathering topsoil samples from several plots). (*) is the point where $y = \delta^{13}\text{C}$ value of the mixture of 63% forest organic matter and 37% savanna organic matter, and corresponding $x=D$ as defined in the text

transect (savanna, edge, forest), by gathering topsoil samples (0–5 cm) from several points (Fig. 5).

In the case of replacement of a savanna vegetation by forest vegetation, the $\delta^{13}\text{C}$ value of SOM is intermediate between $\delta^{13}\text{C}$ values of typical savanna and forest humus. Comparison of the intermediate value with the MRT as given by the ^{14}C datings provides information about the speed of replacement of the initial vegetation.

Results

Botanical study of the Makaba transect

All isolated enclosed savannas show evidence of forest expansion. In a few cases, the expansion is rapid and the forest-savanna interface is invaded by *Aucoumea klaineana*, as described by Foresta (1990). The studied transect, located near Makaba (Figs. 1, 2, transect number 1) corresponds to the general case of a slow forest expansion.

If burning influence is obvious in the savanna part of the vegetation profile, it is weak in the forest part. Flames reach the forest edge, but do not enter into the understorey. This fact is clear from the lack of traces of burning beyond the first shrubs of the boundary. The profile is presented in Fig. 3; Table 1 lists trees whose DBH (diameter at breast high=1.20 m) is greater than 10 cm, and Table 2 the list of $1 < \text{DBH} < 10$ cm trees.

Forest encroaching on savanna can be proved from two different viewpoints.

Floristic proof

Pioneer trees are abundant at the border (Fig. 3: *Aucoumea klaineana*, no. 59; *Pentaclethra eetveldeana*, nos. 22, 48, 50, 54, 56; *Parkia bicolor*, no. 52; *Hymenocardia ulmoides*, no. 62, *Uapaca guineensis*, nos. 46, 58, 61; *Croton* sp., no. 60). Tree species of primary forest only appear in the inner part (*Strombosiopsis tetrandra*, no. 23; *Coula edulis*, nos. 5, 8; *Dacryodes* cf. *igaganga*, nos. 4, 12, 21; *Garcinia* sp., no. 9; *Monodora* cf. *tenuifolia* no. 3; *Dialium* spp., nos. 17, 31). No savanna plants can be found under the forest cover. Outside the forest, some shrubs, too small to be represented in Fig. 3, are new anchor points for the forest expansion. These shrubs are almost always *H. ulmoides* and *Pentaclethra eetveldeana*.

Structural proof

Trees and shrubs which are growing at the border lean towards the savanna. This is due to uneven light conditions during growth, and shows that these trees developed on the edge of the forest and not in the understorey. Other trees (*Parkia bicolor*, nos. 38 and 52), which are now far inside the forest, also have this growth habit, which implies that they once grew at the edge. The morphology of these trees is a precious witness to the forest expansion.

The clearness of the transition between the savanna and forest, vegetation characterized by a "curtain" of almost-vertical leaf surfaces stretched out between the ground and the forest canopy, shows that the expansion of the forest onto the savanna is in this case very slow. If it was rapid, the forest-savanna interface would be less clear, with the presence of grasses and ferns under the trees, similar to some zones invaded by *Aucoumea* already described by Foresta (1990).

Table 1 Transect no. 1. List of diameter at breast height (DBH) >10 cm trees (20×50=1000 m²; see Fig. 3)

Species	Species
1 <i>Aidia micrantha</i> (K. Schum.) F. White	34 <i>Xylopia hypolampra</i> Mildbread
2 <i>Sapindaceae</i> sp. cf. <i>Pancovia</i> ?	35 <i>Dyospyros</i> sp.
3 <i>Monodora</i> cf. <i>tenuifolia</i> Benth	36 <i>Dacryodes</i> cf. <i>igaganga</i> Aubr. & Pellegr.
4 <i>Dacryodes</i> cf. <i>igaganga</i> Aubr. & Pellegr.	37 <i>Aidia micrantha</i> (K. Schum.) F. White
5 <i>Coula edulis</i> Baill.	38 <i>Parkia bicolor</i> A. Cheval.
8 <i>Coula edulis</i> Baill.	39 <i>Pausynistalia</i> sp. 1
9 <i>Garcinia</i> sp.	40 <i>Santiria trimera</i> (Oliv.) Aubr.
10 ?	41 <i>Pausynistalia</i> sp. 1
11 <i>Plagiostyles africana</i> Prain. ex De Wild.	42 <i>Pausynistalia</i> sp. 1
12 <i>Dacryodes</i> cf. <i>igaganga</i> Aubr. & Pellegr.	43 <i>Coelocaryon preussii</i> Warb.
13 <i>Parkia bicolor</i> A. Cheval.	44 <i>Hymenocardia ulmoides</i> Oliver
14 ?	46 <i>Uapaca guineensis</i> Muell. Arg.
15 <i>Parkia bicolor</i> A. Cheval.	47 cf. <i>Duboscia</i> ?
16 Olacaceae sp.	48 <i>Pentaclethra eetveldeana</i> De Wild. et Th. L.
17 <i>Dialium</i> sp.	49 <i>Parkia bicolor</i> A. Cheval.
18 Rubiaceae sp.?	50 <i>Pentaclethra eetveldeana</i> De Wild. et Th. L.
19 ?	51 <i>Santiria trimera</i> (Oliv.) Aubr.
20 <i>Psychotria</i> cf. <i>venosa</i> (Hiern) Petit	52 <i>Parkia bicolor</i> A. Cheval.
21 <i>Dacryodes</i> cf. <i>igaganga</i> Aubr. & Pellegr.	53 <i>Santiria trimera</i> (Oliv.) Aubr.
22 <i>Pentaclethra eetveldeana</i> De Wild. et Th. L.	54 <i>Pentaclethra eetveldeana</i> De Wild. et Th. L.
23 <i>Strombosiopsis tetrandra</i> Engl.	55 <i>Plagiostyles africana</i> Prain. ex De Wild.
24 <i>Symphonia globulifera</i> L. f.	56 <i>Pentaclethra eetveldeana</i> De Wild. et Th. L.
25 <i>Pausynistalia</i> sp. 1	57 <i>Santiria trimera</i> (Oliv.) Aubr.
26 <i>Uapaca</i> sp.	58 <i>Croton</i> sp.
27 <i>Symphonia globulifera</i> L. f.	59 <i>Aucoumea klaineana</i> Pierre
28 cf. Tiliaceae?	60 <i>Croton</i> sp.
29 <i>Parkia bicolor</i> A. Cheval.	61 <i>Uapaca guineensis</i> Muell. Arg.
30 ?	62 <i>Hymenocardia ulmoides</i> Oliver
31 <i>Dialium</i> sp.	6 ? climber
33 <i>Aidia micrantha</i> (K. Schum.) F. White	7 ? climber

Pedological field observations

Soils under savanna-forest transects are of the same kind. Parent rocks, weathering horizons and stone-lines are absolutely similar. Differences only appear in the upper horizons.

The thickness of humus layers ranges from 3 to 7 cm under forest. In savannas, these horizons range from 15 to 25 cm corresponding to the depth of the root system, and extend into the transition horizon A/B. These differences are linked with the humification processes. In forest, the main part of SOM originates from leaves and branches which decay on the soil surface. In savanna, aerial organs of Graminaeae burn every year and SOM are only formed from root exudation and from the decomposition of a dense root system. Therefore, the differences in vegetation are not the result of soil variations. In other word, the few differences between forest and savanna soils result from variation in vegetation.

Erosion steps are very common in savanna. They generally stop at the savanna-forest boundary. Some continue a little under forest, suggesting a great stability of this interface or that forest is slowly encroaching onto savanna. Widely extended erosion steps into forest can only be observed in *Aucoumea* forests (Foresta 1990). This case provides formal proof of the rapid expansion of forest vegetation.

Organic carbon contents

The variations of the soil organic carbon content in the 0–5 cm horizon (Fig. 4) are similar in the two transects. Carbon contents are constant from the savanna up to about 12 m inside the forest. At this stage, the carbon content increases significantly, but very unequally. This irregular increase can be related to the great spatial heterogeneity of soil characteristics in all forests of the Mayombe area at the scale of some square metres (Tsakala 1988; D. Schwartz, unpublished data).

$\delta^{13}\text{C}$ values

The variations of $\delta^{13}\text{C}$ in the 0–5 cm horizon are similar in the two transects (Fig. 5). The $\delta^{13}\text{C}$ is constant under savanna and decreases under and beyond the fern zone. However, it becomes typical of forest only at 17 m (Makaba-transect), and 31 m (Kwilila transect) from the very abrupt Gramineae-fern limit, or at 12 m inside forest. These two points correspond exactly to the increase in organic carbon.

^{14}C ages of SOM

^{14}C ages of SOM (Table 3) are reported in Fig. 5. In the savanna part of the transect, the MRT of SOM in humus layer is 88 years for Kwilila and 64 years for Makaba.

Table 2 Transect no. 1. List of <1 cm <DBH <10 cm individuals (5×50=250 m²)

Species	Species
1 <i>Dialium</i> sp. 1	95 <i>Uapaca guineensis</i> Muell. Arg.
2 Sapindaceae sp. cf. <i>Pancovia</i>	97 cf. <i>Acioa</i> sp.
3 Sapindaceae sp. cf. <i>Pancovia</i>	98 <i>Aidia micrantha</i> (K. Schum.) F. White
4 <i>Dialium</i> sp. 2	99 cf. Chrysobalanaceae sp. 1
5 Sapindaceae sp. cf. <i>Pancovia</i>	110 ?
6 Sapindaceae sp. cf. <i>Pancovia</i>	113 ?
7 <i>Diospyros</i> sp.	114 ?
8 Sapindaceae sp. cf. <i>Pancovia</i>	115 <i>Pentaclethra eetveldeana</i> De Wild. et Th. Dur.
9 <i>Aidia micrantha</i> (K. Schum.) F. White	116 <i>Pentaclethra macrophylla</i> Benth
10 cf. <i>Acioa</i> sp.	117 ?
11 cf. <i>Enantia</i> sp.	118 cf. <i>Duboscia</i> sp.
12 <i>Dialium</i> sp. 3	119 <i>Dialium</i> sp. 2
13 <i>Dialium</i> sp. 4	121 <i>Pentaclethra eetveldeana</i> De Wild. et Th. Dur.
14 Sapindaceae sp. cf. <i>Pancovia</i>	122 ?
15 <i>Carapa procera</i> D.C.	123 cf. Chrysobalanaceae sp. 1
16 <i>Oxyanthus schumannianus</i> De Wild. et Th. Dur.	125 cf. Tiliaceae
17 Chrysobalanaceae sp.	126 <i>Hymenocardia ulmoides</i> Oliver
18 Sapindaceae sp. cf. <i>Pancovia</i>	127 <i>Aidia micrantha</i> (K. Schum.) F. White
19 <i>Dacryodes</i> sp.	128 ?
20 Annonaceae sp.	129 <i>Hymenocardia ulmoides</i> Oliver
21 <i>Dialium</i> sp. 5	130 Meliaceae sp. 2
22 cf. <i>Acioa</i> sp.	131 Meliaceae sp. 2
24 Sapindaceae sp. cf. <i>Pancovia</i>	132 <i>Hymenocardia ulmoides</i> Oliver
26 Sapindaceae sp. cf. <i>Pancovia</i>	133 <i>Hymenocardia ulmoides</i> Oliver
28 cf. Chrysobalanaceae sp. 1	134 <i>Hymenocardia ulmoides</i> Oliver
30 cf. <i>Enantia</i> sp.	135 <i>Hymenocardia ulmoides</i> Oliver
32 cf. <i>Acioa</i> sp.	136 <i>Hymenocardia ulmoides</i> Oliver
33 <i>Dialium</i> sp. 2	23 Climber?
34 <i>Psychotria</i> cf. <i>venosa</i> (Hiern.) E. Petit	25 Connaraceae sp. 1 (climber)
35 Sapindaceae sp. cf. <i>Pancovia</i>	27 Hippocrateaceae sp. 1 (climber)
37 <i>Dacryodes</i> sp.	29 <i>Salacia debilis</i> (G. Don.) Walp. (climber)
38 cf. Chrysobalanaceae sp. 1	31 Hippocrateae sp. 2 (climber)
41 Sapindaceae sp. cf. <i>Pancovia</i>	36 Climber
46 cf. <i>Dialium</i> sp.	39 <i>Salacia debilis</i> (G. Don.) Walp. (climber)
47 Sapindaceae sp. cf. <i>Pancovia</i>	40 <i>Salacia debilis</i> (G. Don.) Walp. (climber)
48 <i>Aucoumea klaineana</i> Pierre	42 Climber
49 cf. <i>Duboscia</i> sp.	43 <i>Salacia debilis</i> (G. Don.) Walp. (climber)
50 Chrysobalanaceae sp.	44 <i>Salacia debilis</i> (G. Don.) Walp. (climber)
52 cf. Chrysobalanaceae sp. 1	45 <i>Artabotrys</i> sp. (climber)
53 <i>Aidia micrantha</i> (K. Schum.) F. White	51 Connaraceae sp. 1 (climber)
54 cf. <i>Acioa</i> sp.	56 Connaraceae sp. 1 (climber)
55 <i>Trichilia</i> sp.	67 Connaraceae sp. 1 (climber)
57 cf. <i>Acioa</i> sp.	68 Climber
58 Sapindaceae sp. cf. <i>Pancovia</i>	70 Connaraceae sp. 1 (climber)
59 Sapindaceae sp. cf. <i>Pancovia</i>	71 Apocynaceae sp. (climber)
60 <i>Uapaca guineensis</i> Muell. Arg.	72 <i>Salacia debilis</i> (G. Don.) Walp. (climber)
61 <i>Symphonia globulifera</i> L.f.	74 Connaraceae sp. 1 (climber)
62 <i>Santiria trimera</i> (Oliv.) Aubr.	75 <i>Salacia debilis</i> (G. Don.) Walp. (climber)
63 <i>Pentaclethra macrophylla</i> Benth.	82 Apocynaceae sp. (climber)
64 <i>Parkia bicolor</i> A. Cheval.	86 Connaraceae sp. 1 (climber)
65 <i>Parkia bicolor</i> A. Cheval.	93 Connaraceae sp. 2 (climber)
66 cf. Chrysobalanaceae sp. 1	94 Connaraceae sp. 1 (climber)
69 cf. Chrysobalanaceae sp. 1	96 Connaraceae sp. 3 (climber)
73 cf. Chrysobalanaceae sp. 1	100 Connaraceae sp. 1 (climber)
77 <i>Uapaca guineensis</i> Muell. Arg.	101 Connaraceae sp. 1 (climber)
78 <i>Dialium</i> sp. 2	102 Connaraceae sp. 1 (climber)
79 Rubiaceae sp.	103 Connaraceae sp. 1 (climber)
80 Meliaceae sp. 1	104 Connaraceae sp. 1 (climber)
83 <i>Dialium</i> sp. 2	105 Connaraceae sp. 1 (climber)
84 cf. Chrysobalanaceae sp. 1	106 Connaraceae sp. 1 (climber)
85 Meliaceae sp. 2	107 Climber
87 Rubiaceae sp.	108 Connaraceae sp. 1 (climber)
88 cf. Euphorbiaceae	109 Connaraceae sp. 1 (climber)
89 cf. Chrysobalanaceae sp. 1	111 Connaraceae sp. 2 (climber)
90 Meliaceae sp. 2	112 Connaraceae sp. 1 (climber)
91 <i>Pentaclethra eetveldeana</i> De Wild. et Th. Dur.	120 Climber
92 <i>Oxyanthus schumannianus</i> De Wild. et Th. Dur.	124 Connaraceae sp. 3 (climber)

Table 3 ^{14}C activity in year 1990 and mean residence time (MRT) of soil organic matter of average samples in different parts of the two transects

Sample	$\delta^{14}\text{C}$ (‰)	$\delta^{13}\text{C}$ (‰)	$\Delta^{14}\text{C}$ (‰)	MRT (years)
Makaba, savanna	171.7	-12.5	142.4	64
Makaba, shrub	180.9	-24.3	179.2	45
Makaba, forest	213.9	-27.0	218.8	32
Kwilila, savanna	138.8	-14.0	113.7	88
Kwilila, fern	122.0	-21.0	113.0	88
Kwilila, forest	210.6	-27.0	215.4	32

MRT of the sample from the shrub-zone and forest edge at Kwilila is close to that in the savanna. MRT of humus at the forest edge in Makaba is 45 years. In the forest, the MRT is 32 years in both cases. The lower MRT of SOM in the forest, when compared to the savanna, can be related to the higher carbon content. It can be explained by an increased carbon input to the soil surface and/or a higher turnover rate.

Discussion

Proof of forest encroachment

MRT and $\delta^{13}\text{C}$ indicate changes of biogeochemical characteristics of SOM. Are these facts proof of forest expansion or are they the observation of an edge effect occurring on a static boundary?

An answer is given according to recent data in Amazonia by Kapos et al. (1993), suggesting that the $\delta^{13}\text{C}$ composition of canopy leaves, understorey leaves and understorey air changes in relation with the distance to the edge in a 5-year-old static edge with frequent gaps. However, the response of two understorey species studied is different. The $\delta^{13}\text{C}$ composition of leaves of the

first species does not present significant changes between edge and forest zones, whereas the $\delta^{13}\text{C}$ in leaves of the second species presents a decrease of 2 ‰ in absolute value from the edge up to 40 m in the forest. As a general rule, Kapos et al. (1993) never found decreases greater than 2 ‰ as edge effects.

On the other hand three different arguments lead us to consider that in the transects forest is encroaching on savanna:


Floristic and structural proof of forest expansion

The present position of trees showing ripicole behaviour and increasing diameters in relation with their distance to the edge, such as *Parkia bicolor* trees no. 57 (DBH=27 cm; distance from the edge: 6 m) and no. 38 (DBH=43 cm; distance: 10 m), provide a reference point as they germinated at the forest boundary.

$\delta^{13}\text{C}$ of SOM

Even if the 2 ‰ differences in the $\delta^{13}\text{C}$ values of leaves and atmospheric CO_2 , observed by Kapos et al. (1993) could be transferred into litterfall, litter composition on the soil and into the following $\delta^{13}\text{C}$ values of SOM in humus layer, no edge effect can explain differences as high as 14 ‰. Moreover, the values of the $\delta^{13}\text{C}$ of SOM in forest humus layers at 12–17 m from the edge in both transects are becoming identical to the $\delta^{13}\text{C}$ values of other typical rain-forests from the south of the Congo, which range from -26.6 to -28.2 ‰ (Schwartz 1991). In other words, if the ecotone was static, differences between the $\delta^{13}\text{C}$ of SOM in savanna and forest topsoils should be more abrupt, in spite of the edge effects, as schematized in Fig. 6. Moreover the C4–C3 edge is not the savanna-forest ecotone, but the graminean-fern limit.

Fig. 6A–C Theoretical $\delta^{13}\text{C}$ variations in a savanna-forest transect in the three following hypotheses: A static ecotone, B savanna expansion, C forest encroachment. In a simplified manner: C4=graminean zone; C3=fern patch+forest. NB The inclination of the slope is an indicator of the speed of the encroachment



This fact could reduce the edge effect, which is a straight consequence of the difference of the $\delta^{13}\text{C}$ composition between atmospheric and understorey CO_2 .

Organic carbon under forest

As additional proof, the increase in organic carbon content under forest also shows that this vegetation type is progressing compared to savanna in the two transects. If the savanna-forest boundary was stable, the carbon increase should only appear just beyond the boundary. If savanna is progressing, the carbon content under savanna near the edge should be comparable to the carbon content under forest. The irregular increase of organic carbon content is to be referred with the great lateral variability of topsoil horizons in the Mayombe forests on a metric scale (Tsakala 1988).

Estimation of the speed of forest progression

Combined $\Delta^{14}\text{C}$ and $\delta^{13}\text{C}$ data allowed us to estimate the speed of forest encroachment. According to the model of exponential decay we used to compute MRT of SOM after $\Delta^{14}\text{C}$ values in top soil samples (Balesdent and Guillet 1982), MRT is the time required for the replacement of 63% of the savanna SOM by forest SOM [i.e. $1 - \exp(-1)$]. We measured the distance D between typical savanna $\delta^{13}\text{C}$ values and $\delta^{13}\text{C}$ values corresponding to the mixture of 63% forest organic matter and 37% savanna organic matter. Then we estimated that the distance D was covered within the time MRT:

$$\text{Speed} = D \cdot \text{MRT}^{-1} \text{ m/century.}$$

We estimated D from the data in Fig. 5. At Makaba (transect number 1) it is equal to 9 m. As MRT ranges between 64 and 32 years, the present speed of forest encroachment is between 14 and 28 m/century. At Kwilila (transect number 2), the same calculation, with $D=24$ m and $\text{MRT}=32-88$ years, leads to a speed between 27 and 75 m/century. As MRT progressively decreases from the savanna to the forest, medium values may be a better approximation of forest progression. We finally estimate that the mean speed of forest encroachment in enclosed savannas of the Mayombe probably ranges between 20 and 50 m century. This speed is slow in the absolute. Annual burnings of savannas explain this fact, but it is important to notice that burning cannot prevent forest progression.

History of vegetation

The history of the vegetation in the south of the Congo was reconstructed by palynological data, vegetable macroremains determination and $\delta^{13}\text{C}$ analyses (Dechamps et al. 1988; Elenga et al. 1992; Schwartz 1988, 1991, 1992; Schwarz et al. 1992; Vincens et al. 1994). These

data show that forest covered all this area during early Holocene, that a climatic drying at about 3000 B.P. led to the expansion of the present savannas and that the climate became more humid again some centuries ago. The present forest encroachment in the enclosed savannas could correspond to this climatic episode. As a consequence, the savannas were certainly more extensive some centuries ago, and this confirms some archaeological data of Schwartz et al. (1990a), according to which savannas occurred within the Mayombean forest as far west as Les Saras.

Conclusions

The botanical study, organic carbon contents and $\delta^{13}\text{C}$ values on the two transects show undoubtedly that forest is encroaching onto savanna. The climatic conditions within the Mayombe are favourable to forest expansion. This expansion can be seen in the upper horizon. The transformation starts in the fern zone; ferns are C3 plants. Combination of $\Delta^{14}\text{C}$ and $\delta^{13}\text{C}$ data allows one to estimate that the speed of expansion in the two representative transects is between 14 and 75 m/century, and more likely at about 20–50 m/century. It is clear that savanna bush fires slow down the forest encroachment, but it is also clear that they cannot prevent it. As most savannas are less than 1 km across, speeds of 20–50 m/century mean that, assuming environmental conditions remain the same, these savannas could disappear in about 1 or 2 millennia. The area of about 40% of the savannas is smaller than 0.1 km² (Rat Patron and Schwartz 1996); these little savannas could disappear in 100–200 years.

This expansion speed also leads to questions about the regeneration speed of upper horizons of soil. As opposed to savanna, the latter are never scoured down to the stone-line under forest. It seems very possible that these horizons can be regenerated under forest by means of different factors which remain to be precisely studied (biological factors, especially termite intervention and accumulation of colluviated particles). This regeneration could occur in a few centuries, which is very rapid in pedological terms.

Acknowledgements The research was supported by the French CNRS and ORSTOM through the International Program ECOFIT (Ecologie et Paléocologie des forêts intertropicales – Ecology and Paleocology of the Intertropical Forests).

References

- Balesdent J, Guillet B (1982) Les datations par le ^{14}C des matières organiques du sol. Contribution à l'étude de l'humification et du renouvellement des substances organiques. *Sci Sol* 2:93–112
- Balesdent J, Mariotti A, Guillet B (1987) Natural ^{13}C abundance as a tracer for studies of soil organic matter dynamics. *Soil Biol Biochem* 19:25–30
- Bender MM (1971) Variations in the $^{13}\text{C}/^{12}\text{C}$ ratios of plants in relation to the pathway of photosynthetic carbon dioxide fixation. *Phytochemistry* 10:1239–1244

- Benner R, Fogel ML, Sprague EK, Hodson RE (1987) Depletion in ^{13}C lignin and its implications for stable isotope studies. *Nature* 329:708–710
- Blair N, Leu A, Munoz E, Olsen J, Kwong E, Des Marais D (1985) Carbon isotope fractionation in heterotrophic microbial metabolism. *Appl Environ Microbiol* 50:996–1001
- Clairac B, Cros B, Senechal J (1989) Le climat du Mayombe. In: *Revue des connaissances sur le Mayombe*. PNUD, MAB, UNESCO, Paris, pp 47–68
- Cusset G (1989) La flore et la végétation du Mayombe congolais. Etat des connaissances. In: *Revue des connaissances sur le Mayombe*. PNUD, MAB, UNESCO, Paris, pp 103–136
- Dechamps R, Guillet B, Schwartz D (1988) Découverte d'une flore forestière mi-Holocène (5800–3100 B.P.) conservée in situ sur le littoral ponténégrin (R.P. du Congo). *CR Acad Sci Paris Ser II* 306:615–618
- Dzurec RS, Boutton TW, Caldwell MM, Smith BN (1985) Carbon isotope ratios of soil organic matter and their use in assessing community composition change in Curlew Valley, Utah. *Oecologia* 66:17–24
- Elenga H, Schwartz D, Vincens A (1992) Changements climatiques et action anthropique sur le littoral congolais au cours de l'Holocène. *Bull Soc Geol Fr* 163:83–90
- FAO (1976) Carte mondiale des sols, vol VI. Afrique. FAO, Rome
- Flexor JM, Volkoff B (1977) Distribution de l'isotope stable ^{13}C dans la matière organique d'un sol ferrallitique de l'Etat de Bahia (Brésil). *CR Acad Sci Paris Ser D* 284:1655–1657
- Foresta H de (1990) Origine et évolution des savanes intramayombiennes (RP du Congo). II. Apports de la botanique forestière. In: Lanfranchi R, Schwartz D (eds) *Paysages quaternaires de l'Afrique Centrale atlantique*. ORSTOM, Paris, pp 326–335
- Gibert G, Senechal J (1989) L'économie forestière. In: *Revue des connaissances sur le Mayombe*. PNUD, MAB, UNESCO, Paris, pp 249–293
- Girardin C, Mariotti A (1991) Analyse isotopique du ^{13}C en abondance naturelle dans le carbone organique: un système automatique avec robot préparateur. *Cah ORSTOM Ser Pedol* 26:371–380
- Goh KM, Rafter TA, Stout JD, Walker TW (1976) The accumulation of soil organic matter and its carbon isotope content in a chronosequence of soils developed on aeolian sand in New Zealand. *J Soil Sci* 27:89–100
- Goh KM, Stout JD, Rafter TA (1977) Radiocarbon enrichment of soil organic matter fractions in New Zealand soils. *Soil Sci* 123:385–391
- Gras F (1970) Surfaces d'aplanissement et remaniement des sols sur la bordure orientale du Mayombe. *Cah ORSTOM Ser Pedol* 8:274–294
- Kaplan IR, Rittenberg SC (1964) Carbon isotope fractionation during metabolism of lactate by *Desulfovibrio desulfuricans*. *J Gen Microbiol* 34:213–217
- Kapos V, Ganade G, Matsui E, Victoria RL (1993) $\delta^{13}\text{C}$ as an indicator of edge effects in tropical rainforest reserves. *J Ecol* 81:425–432
- Marino BD, McElroy MB (1991) Isotopic composition of atmospheric CO_2 inferred from carbon in C4 plant cellulose. *Nature* 349:127–131
- Mariotti A (1991) Le carbone 13 en abondance naturelle, traceur de la dynamique de la matière organique des sols et de l'évolution des paléoenvironnements continentaux. *Cah ORSTOM Ser Pedol* 26:299–313
- Mariotti A, Balesdent J (1990) ^{13}C natural abundance as a tracer of soil organic matter turnover and paleoenvironment dynamics. *Chem Geol* 84:217–219
- Mariotti A, Peterschmitt E (1994) Forest savanna ecotone dynamics in India as revealed by carbon isotope ratios of soil organic matter. *Oecologia* 97:475–480
- Martin A, Mariotti A, Balesdent J, Lavelle P, Vuattoux R (1990) Estimate of organic matter turn-over rate in a savanna soil by ^{13}C natural abundance measurements. *Soil Biol Biochem* 22:517–523
- Massimba JP (1987) Etude descriptive d'une forêt à Marantaceae/Zingiberaceae dans le Mayombe congolais. DEA, Université de Paris VI
- Nadelhoffer KJ, Fry B (1988) Controls on natural nitrogen-15 and carbon-13 abundances in forest soil organic matter. *Soil Sci Soc Am J* 52:1633–1640
- Nissenbaum A, Schallinger KM (1974) The distribution of the stable carbon isotope ($^{13}\text{C}/^{12}\text{C}$) in fractions of soil organic matter. *Geoderma* 11:137–145
- O'Brien BJ, Stout JD (1978) Movement and turnover of soil organic matter as indicated by carbon isotope measurements. *Soil Biol Biochem* 10:309–317
- Petit M (1990) Géographie physique tropicale. Karthala-ACCT, Paris
- Rat Patron P, Schwartz D (1996) Les savanes incluses du Mayombe. I. Description et localisation. In: Reversat F, Schwartz D (eds) *Milieux et activités agricoles dans le Mayombe congolais*. UNESCO, Paris (in press)
- Schwartz D (1988) Histoire d'un paysage: le lousseke. Paléoenvironnements quaternaires et podzolisation sur sables Bateke (quarante derniers millénaires, région de Brazzaville, RP du Congo). ORSTOM, Paris
- Schwartz D (1991) Intérêt de la mesure du $\delta^{13}\text{C}$ des sols en milieu naturel équatorial pour la connaissance des aspects pédologiques et écologiques des relations savane-forêt. *Cah ORSTOM Ser Pedol* 26:327–341
- Schwartz D (1992) Assèchement climatique vers 3000 BP et expansion bantu en Afrique Centrale atlantique: quelques réflexions. *Bull Soc Geol Fr* 163:353–361
- Schwartz D, Mariotti A, Lanfranchi R, Guillet B (1986) $^{13}\text{C}/^{12}\text{C}$ ratios of soil organic matter as indicators of vegetation changes in the Congo. *Geoderma* 39:97–103
- Schwartz D, Foresta H de, Dechamps R, Lanfranchi R (1990a) Découverte d'un premier site de l'Age du fer ancien (2110 BP) dans le Mayombe congolais. Implications paléobotaniques et pédologiques. *CR Acad Sci Paris Ser II* 310:1293–1298
- Schwartz D, Lanfranchi R, Mariotti A (1990b) Origine et évolution des savanes intramayombiennes (RP du Congo). I. Apports de la pédologie et de la biogéochimie isotopique (^{14}C et ^{13}C). In: Lanfranchi R, Schwartz D (eds) *Paysages quaternaires de l'Afrique Centrale atlantique*. ORSTOM, Paris, pp 314–325
- Schwartz D, Mariotti A, Trouvé C, Van Den Borg K, Guillet B (1992) Etude des profils isotopiques ^{13}C et ^{14}C d'un sol ferrallitique sableux du littoral congolais. Implications sur la dynamique de la matière organique et l'histoire de la végétation. *CR Acad Sci Paris Ser II* 315:1411–1417
- Smith BN, Epstein S (1971) Two categories of $^{13}\text{C}/^{12}\text{C}$ ratios for higher plants. *Plant Physiol* 47:380–384
- Troughton JH, Wells PV, Mooney HA (1974) Photosynthetic mechanisms and paleoecology from carbon isotope ratios in ancient specimens of C4 and CAM plants. *Science* 185:610–612
- Tsakala R (1988) Etude de la variabilité spatiale des caractéristiques analytiques du sol de deux parcelles sous forêt et sous bananiers dans le Mayombe. *Mem Dipl Ing Develop Rural*, ORSTOM Pointe Noire and IDR, Brazzaville
- Vennetier P (1968) Pointe Noire et la façade maritime du Congo. Mémoire no 26. ORSTOM, Paris
- Vincens A, Buchet G, Elenga H, Fournier M, Martin L, Namur C de, Schwartz D, Servant M, Wirrmann D (1994) Un changement majeur de végétation autour du lac Sinnda (vallée du Niari, sud-Congo) lié à un assèchement climatique intra-Holocène. Apport de la palynologie. *CR Acad Sci Paris Ser II* 318:1521–1526
- Volkoff B, Cerri CC (1987) Carbon isotopic fractionation in subtropical Brazilian grassland soils. Comparison with tropical forest soils. *Plant Soil* 102:27–31
- Walkley A, Black A (1934) An examination of the Degtjareff method for determining soil organic matter and a proposed modification of the chronic acid titration method. *Soil Sci* 37:29–38
- White F (1983) The vegetation of Africa. UNESCO/AET-FAT/UNSO, Paris