

MANAGING SOIL FERTILITY ON TERRACES FORMING BEHIND VEGETATIVE FILTER STRIPS: AN ASSESSMENT OF FARMERS' STRATEGIES

M. Stark¹; D.P. Garrity² and S.C. Jutzi¹

¹ *Institute of Crop Science, University of Kassel, Steinstr. 19, 37213 Witzenhausen, Germany*

² *Southeast Asian Regional Research Programme, International Centre for Research in Agroforestry (ICRAF), P.O. Box 161, Bogor 16001, Indonesia*

ABSTRACT

The indigenous use of natural vegetative strips (NVS) to control soil erosion on the slope has been viewed as a low-cost alternative to planted tree hedgerows. As in conventional hedgerow systems, however, natural terrace formation resulting from redistribution of sediment from upper to lower terrace zones leads to the development of a soil fertility gradient with significantly lower crop yield on the degraded upper portion of each terrace. Since NVS produce little biomass which could be used to maintain soil fertility, the sustainability of annual crop production in NVS systems may be questionable. Interview surveys conducted in two upland locations in the Philippines showed that most farmers had observed soil fertility scouring to adversely affect crop performance on the upper part of the terrace. However, scouring was not usually perceived as a serious constraint of the technology. Farmers claimed that the benefits of overall increased crop yield and rise in land value due to contouring outweighed the negative effects of upper terrace yield decline; besides, they generally believed scouring to be a transitory phenomenon. On-farm experiments were conducted to assess farmers' strategies to overcome the negative effects of soil fertility scouring in NVS systems by skewing the application of nutrient inputs towards degraded upper terrace zones. Further research is required to identify methods to fully rehabilitate the degraded upper terrace through raising soil organic matter levels.

1. INTRODUCTION

The indigenous practice of using natural vegetative strips (NVS) of 0.5 - 1 meter width as buffer or filter strips to reduce soil erosion on the slope can be viewed as an alternative technology to contour hedgerow intercropping with planted trees. It requires minimal labor for establishment and maintenance while reducing soil erosion at least as effectively as planted tree hedgerows (Garrity *et al.*, 1993). Local grasses and broad-leaf species compete less with adjacent alley crops (Ramiamanana, 1993). However, sediment redistribution during natural terrace formation leads to the development of a soil fertility gradient with significantly lower crop yield on degraded upper terrace (alley) zones as compared to the lower terrace portion where eroded sediment and nutrients accumulate (Turkelboom *et al.*, 1993; Anecksamphant and Sajjapongse, 1994). Natural vegetative contour strips provide substantially less biomass compared to tree hedgerows which could be used to maintain soil fertility levels on the upper terrace and reduce the effects of scouring. Sustained continuous crop production in NVS systems therefore depends largely on the import of external nutrients.

Research has focused on significantly reducing within-alley soil movement through ridge tillage cultivation, a minimum tillage technology (Thapa *et al.*, 1996), and on biasing the application of hedgerow cuttings and crop residues towards degraded upper alley (terrace) zones in tree hedgerow systems (Mercado *et al.*, 1996). The suggestion to improve fertilizer use efficiency by matching fertilizer rates to site specific nutrient requirements at varied landscape positions (Fiez *et al.*, 1994) may also be applied to small-scale variations of soil fertility in vegetative contour strip systems.

The present study is part of a participatory technology development (PTD) project conducted under ICRAF's research program in the Philippines. Objectives of the study were to identify farmers' indigenous strategies to improve the NVS system with a special focus on overcoming the negative effects of soil fertility scouring on crop yield. Subsequent on-farm research focused on assessing indigenous strategies to obtain uniform crop yield across the alley-way by biasing the application of nutrient inputs towards degraded upper alley zones.

A deeper understanding of the benefits and constraints of natural vegetative contour strips, together with identifying and developing options for the improved management of NVS systems, will allow more confident extrapolation of the technology to other locations in the humid tropics.

2. METHODOLOGY

Fifty-three interviews were conducted among contour farmers in Claveria (Northern Mindanao) and Matalom (Southern Leyte) in 1995 and 1997, respectively. Claveria municipality is one of ICRAF's research sites in the

Philippines located in an acidic upland area typical for the majority of degraded uplands in the SE-Asian region. In Matalom, the use of natural vegetative strips is an indigenous soil conservation technology which has been applied by local farmers for more than 50 years. Matalom is characterized by shallow calcareous soil conditions and less intensive cropping. Basic crop and soil data was collected from selected farms to document the effects of soil fertility scouring. Formal on-farm experiments conducted on existing NVS sites since 1995 assess identified farmer strategies to overcome the effects of soil fertility scouring.

3. RESULTS AND DISCUSSION

3.1 Farmers' perceptions of soil fertility scouring

Most farmers had observed soil fertility scouring to adversely affect plant height and crop yield on the upper part of the alley, but did not consider scouring as a serious constraint of the technology. An example of the dramatic effect of soil fertility scouring on crop yield is shown in Figure 1. Farmers claimed that the benefits of overall increased crop yield and rise in land value due to contouring outweighed the negative effects of upper alley yield decline. In Claveria, scouring was generally observed or believed to be a transitory phenomenon, either affecting crop performance during the first three to five years after NVS establishment or for at least the same duration at a later stage, after more soil had been relocated within alleys. Under less intensive cropping in Matalom, however, it commonly takes several decades until terraces level off and until degraded upper alley soil conditions have been rehabilitated.




Figure 1: Yield of unfertilized native maize and peanut intercrop across a single alley between vegetative contour strips (grain at 14 %, nuts at 12 % moisture content; means of three replications; 1. crop 1995, farmer's practice)

Skewing the application of nutrient inputs towards the degraded upper alley was found to be a common strategy among farmers in Claveria to reduce the negative effects of soil fertility scouring on crop yield (Tab. 1). Since the application of external inputs was not an option for most farmers in Matalom, only few had applied strategies to maintain soil fertility, aside from following the land.

Table 1: Locals strategies to reverse or avoid the effects of soil fertility scouring used among farmers in Claveria (n=30)-

Farmers' Methods to Overcome Scouring Effects	Count	Percent
Up to two times more fertilizer on upper alley crop rows	12	34.3
Applying fertilizer uniformly across alley	5	14.3
Additional mulch application on upper alley crop rows	4	11.4
Skewing the distribution of crop residues towards upper alley	2	5.7
Skewing the application of lime towards upper alley	2	5.7
None	10	28.6
Total*	35	100

* Multiple responses possible

3.2 Technology validation: an assessment of farmers' strategies

On-farm trials to evaluate farmers' management strategies have been conducted in Claveria since 1995. Experiments showed that skewing the application of mineral fertilizer (treatments T3 and T4) towards upper alley zones resulted in significantly higher hybrid maize grain yield on the degraded upper alley compared to uniform fertilizer distribution (T2). The application of 1.5 times more NPK-fertilizer on upper compared to lower alley zones (T3) was optimal and resulted in the lowest crop yield response slope, i.e. the most uniform crop performance across the alley. A three times higher fertilizer application (T4) over-compensated for the within-alley fertility gradient (Fig. 2). NPK fertilizer was applied at the recommended rate of 80-30-30 kg NPK ha⁻¹ season⁻¹ (full fertilizer rate) for three cropping seasons; treatment T1 represents the control with no fertilizer applied. Similar experimental results were obtained at half the recommended fertilizer rate which was tested during the final, i.e. the fourth trial crop.

Looking at grain yield harvested from the alley as a whole, there were no significant differences between uniform and skewed fertilizer treatments, because placing increasingly higher portions of the fertilizer amount on the upper alley constantly reduced yield on the lower alley zones. The yield decline on lower alley zones followed a linear trend, at a rate almost identical with the rate of yield increase on more heavily fertilized upper alley zones. The study was, however, limited to only a small number of treatment levels and maize varieties.

Initial investigation of the skewed distribution of either crop residues or lime showed positive effects on upper alley crop yield. A long-term investigation of methods to fully rehabilitate soil fertility on degraded alley zones through raising soil organic matter levels is required. Computer models could be useful in extrapolating experimental results.




Figure 2: Effect of fertilizer treatments on hybrid maize grain yield across alley zones; 1st crop 1996, full fertilizer rate (Note: LSD values [bars inside graph] computed separately by alley zone for $P < 0.05$)

4. CONCLUSION AND RECOMMENDATIONS

The adoptability of the NVS technology can be enhanced by building on indigenous strategies to improve the NVS system. Further experimental evaluation of farmers' practices is required to identify strategies to fully rehabilitate degraded upper alley soil conditions in the long-run. These strategies are: skewing the application of crop residues, mulch and/or lime with higher amounts on the degraded upper alley zones, and clean-scraping the lower portion of the NVS bunds to add nutrients (with the scraped plant and soil material) to the upper part of the alley.

Other indigenous technology modifications which need to be evaluated before their wider dissemination include:

- Adding value to the area "lost" to NVS and reinforcing buffer function: most farmers perceived NVS as an intermediate step towards the development of more complex agroforestry systems. Interaction effects and overall system productivity need to be evaluated for a variety of perennials planted on the contour strips and a number of different planting arrangements. Under low-intensity cropping, vetiver grass (*Vetiveria zizanioides*) could provide reinforcement for the NVS because low pruning frequency and prolonged fallow periods (with free grazing animals causing damage to the strips) result into weak buffer strips.
- Reducing labor requirements for establishment of NVS: most farmers had substituted labor-intensive methods for simple eye-estimate to determine the (semi)contour line. Loss of efficiency in controlling soil erosion needs to be quantified.

Strengthening the partnership between farmers, researchers and the local government has become an important goal of the research program and proved to be a successful model for the efficient, low-cost and sustained development and dissemination of soil-conserving and income-generating technologies among small-scale farmers in the study area.

5. REFERENCES

Aneksamphant, C. and Sajjapongse, A. (1994): Impacts of soil erosion on crop growth and yield. Paper presented at the International Workshop on Conservation Farming for Sloping Uplands in Southeast Asia: Challenges, Opportunities and Prospects. 20-26 November 1994, Manila, Philippines.

Fiez, T.E., Miller, B.C. and Pan, W.L. (1994): Winter wheat yield and grain protein across varied landscape positions. In: *Agron. J.* 86: 1026-1032.

Garrity, D.P., Kummer, D.M. and Guiang, E.S. (1993): The Philippines. In: National Research Council: Sustainable agriculture and the environment in the humid tropics. National Academy Press, Washington, D.C.: 549-624.

Mercado, A.R., Sanchez, N. and Garrity, D.P. (1996): Crop productivity using forage legumes and grasses as contour hedgerow species in an acid upland soil. Paper presented at the 12th Annual Scientific Conference of the Federation of Crop Sciences Societies of the Philippines, May 13-18, 1996, Davao, Philippines.

Ramiaramanana, D.M. (1993): Crop-hedgerow interactions with natural vegetative filter strips on sloping acidic land. MSc. thesis, University of the Philippines at Los Baños, Philippines.

Thapa, B., Cassel, D.K., Garrity, D.P. and Mercado, A.R. (1996): Sustainable soil management systems on acid steepland soils in the humid tropics. Paper presented at the 12th Annual Scientific Conference of the Federation of Crop Sciences Societies of the Philippines, May 13-18, 1996, Davao, Philippines.

Turkelboom, F., Ongprasert, S. and Taejjai, U. (1993): Alley Cropping on Steep Slopes: Soil Fertility Gradients and Sustainability. Paper presented at the International Workshop on 'Sustainable Agricultural Development: Concepts and Measures', held at AIT, Bangkok, 14-17 December 1993, Bangkok, Thailand.

6. ACKNOWLEDGMENT

The study was supported by the International Centre for Research in Agroforestry (ICRAF) and the *Studienstiftung des Deutschen Volkes*. Research was conducted in the context of the *Alternatives to Slash-and-Burn* (ASB) project, a global and multi-institutional initiative coordinated by ICRAF.

- An Assessment of Local Strategies -

Marco Stark¹, Dennis P. Garrity² and Samuel C. Jutz¹

¹University of Kassel • Institute of Crop Science • Germany • and
²International Centre for Research in Agroforestry (ICRAF) • Philippines



Introduction

The indigenous use of natural vegetative contour strips (NVS) in the Philippine uplands has been viewed as a low-cost alternative to planted tree hedgerows (Plate 1). As in conventional hedgerow systems, however, natural terrace formation resulting from the redistribution of sediment from upper to lower terrace zones (Plate 2) leads to the development of a soil fertility gradient with significantly lower crop yield on the degraded upper portion of each terrace. To sustain annual crop production in NVS systems the application of external nutrients may be required.



Plate 1&2: NVS are established by simply leaving narrow contour strips unplowed during land preparation; native grass and broad-leaved species provide an effective barrier to eroded sediment, leading to rapid terrace formation.

Major Objectives of the study were to identify farmers' strategies to improve the NVS system with a special focus on overcoming the negative effects of soil fertility scouring on crop yield. An understanding of the benefits and constraints of natural vegetative contour strips, together with identifying options for the improved management of NVS systems, will allow more confident extrapolation of the technology to other locations in the humid tropics.

Methodology

Fifty-three interviews were conducted among contour farmers in Claveria (Northern Mindanao) and Matalom (Southern Leyte) in 1995 and 1997, respectively. The municipality of Claveria is one of ICRAF's research sites in the Philippines located in an acidic upland area. Matalom is characterized by shallow calcareous soil conditions and less intensive cropping.

Results and Discussion

Most farmers had observed soil fertility scouring to adversely affect plant height and crop yield on the upper part of the alley (= terrace) between the vegetative contour strips (Fig. 1), but did not consider scouring as a serious constraint of the technology. Farmers claimed that the benefits of overall increased crop yield and rise in land value due to contouring outweighed the negative effects of upper alley yield decline.

In Claveria, scouring was generally observed or believed to be a transitory phenomenon. Under less intensive cropping in Matalom, however, it may take several decades until terraces level off and degraded upper alley soil conditions have been rehabilitated.




Figure 1: Yield of unfertilized native maize and peanut intercrop across a single alley between vegetative contour strips (means of 3 replications; farmer's practice)

The study concluded that the adoptability of the NVS technology can be further enhanced by building on farmers' own strategies to reduce the effects of soil fertility scouring on crop yield. Skewing the application of nutrient inputs towards the degraded upper alley was found to be a common practice among farmers in Claveria (Tab. 1).

Table 1: Local strategies to reverse or avoid the effects of soil fertility scouring used among farmers in Claveria (n=30)

Methods To Overcome Scouring Effects	Count	Percent
Up to two times more fertilizer on upper alley crop rows	12	34.3
Fertilizer application uniformly across alley	5	14.3
Additional mulch application on upper alley crop rows	4	11.4
Skewing crop residues towards upper alley	2	5.7
Skewing the application of lime towards upper alley	2	5.7
None	10	28.6
Total (multiple responses possible)	35	100

Other indigenous technology modifications which need to be evaluated before their wider dissemination, include:

- Adding value to the NVS and reinforcing buffer function: Interaction effects and overall system productivity need to be evaluated for a variety of cash perennials planted on the contour strips. Under low-intensity cropping, vetiver grass (*Vetiveria zizanioides*) could be used to reinforce weak buffer strips.
- Reducing labor requirements for establishment of NVS: the effect of using simple eye-estimate instead of more labor-intensive methods to determine the (semi-) contour line needs to be assessed with regards to labor cost and efficiency in controlling soil erosion.

On-farm trials to evaluate indigenous management strategies have been conducted in Claveria since 1995. Skewing the application of mineral fertilizer inputs towards degraded upper alley zones increased crop yield on this alley portion significantly. However, fertilizer efficiency for the alley as a whole remained unchanged for the fertilizer rates tested. Initial investigation of the skewed distribution of either crop residues or lime showed positive effects on upper alley crop yield. Further investigation of methods to fully rehabilitate soil fertility on degraded alley zones through raising soil organic matter levels are required.

The participatory technology development & dissemination approach employed in the present study proved to be a successful model for efficient, low-cost and sustained research and development.



NATURAL VEGETATIVE STRIPS (NVS)

- A Low-Cost Soil Conservation Technology -

Marco Stark¹, Dennis P. Garrity² and Samuel C. Jutzi¹

¹University of Kassel • Institute of Crop Science • Germany • and
²International Centre for Research in Agroforestry (ICRAF) • Philippines



Farmer-modified Soil Conservation Technology as a Base for Participatory Research

During the 1980s, upland farmers exposed to the concept of contour hedgerow intercropping in Claveria, Northern Mindanao, started experimenting with natural vegetative filter strips (NVS) as a low-cost alternative to introduced tree & grass hedgerows.




The advantages of natural vegetative strips in comparison to planted tree & grass hedgerows are:

- Low labor requirements for establishment and maintenance. Filter strips are simply laid out by leaving 50 cm wide strips along the contour uncultivated which vegetate naturally with local grass and broad-leaved species. Many farmers estimate the contour line by simple eye-estimate (small photo).
- Effective reduction of soil loss: research has shown that NVS reduce soil loss on a 25 % slope typically by more than 90 %.
- Minimal competition effects on adjacent crops.



Natural terrace formation is rapid due to frequent animal tillage and water-induced erosion. Ten years after establishment, these terraces between napier grass strips on a 40 % slope have almost leveled off. Land preparation on the sloping field has become easy.

Rapid soil redistribution during natural terrace development leads to the formation of a soil fertility gradient across individual terraces. The negative effects of soil fertility scouring on crop yield on upper terrace zones can be dramatic.



Farmers' indigenous strategies to deal with soil fertility scouring are being investigated by ICRAF in Claveria. On-farm trials focus mainly on skewing nutrient inputs towards degraded upper terrace zones. Other research includes studying the effects of perennials, planted on the NVS, on alley crop and over-all system productivity.

